

Benefits of using Community Health - Health Promotion resources to act to reduce tobacco-related harm

Your Community Health Service should prioritise community-focused action to reduce tobacco-related harm to:

- ✓ Reduce the **substantial burden** on **healthcare** and the **community** from the wide-ranging morbidity and mortality caused by smoking
- ✓ Achieve your purpose of **prioritising those** with, or **at risk of**, **poorer health**
- ✓ **Reduce** the **COVID-19 risks** and **consequences** exacerbated by smoking and vaping
- ✓ Act early on **concerns** that the increasing use of **e-cigarettes** will lead to **increases in smoking** in young people
- ✓ **Improve** mental health and wellbeing
- ✓ **Tackle poverty** in applying the social model of health
- ✓ Reduce exposure to the significant **adverse effects** of **second- and third-hand smoke** on non-smokers and children
- ✓ Facilitate and strengthen **collaborative** local and state-wide **efforts** to tackle tobacco-related harm
- ✓ **Benefit** from the **support, guidance** and **opportunities** offered by Quit Victoria

Why your Community Health Service should prioritise community-focused action to reduce tobacco-related harm

- ✓ Reduce the substantial burden on healthcare and the community from the wide-ranging morbidity and mortality caused by smoking¹

Smoking kills around
20,000
Australians
each year.



That's **54**
preventable
deaths every day.

Smoking
leads to at least
46
Medical
conditions
requiring
treatment
and care.



Inpatient hospital
treatment
\$1.5 billion



Primary healthcare
treatment
\$1.5 billion



Ambulance and
ED treatment
\$453 million



Pharmacy
medications
\$451 million



Smoking cessation
medications
\$153 million



Outpatient
treatment
\$289 million



Residential care
\$420 million



Family care
\$2.0 billion

The cost of healthcare due to smoking is
\$6.8 billion.

Australia: July 2015 to June 2016

Costliest adverse health outcomes caused by smoking:

\$347.3 million

Chronic obstructive
pulmonary disease

\$205.9 million

Ischaemic heart
disease

\$152.8 million

Tracheal, bronchus
and lung cancer

Other adverse health outcomes caused by smoking:

Heart disease and stroke

- Ischaemic strokes
- Haemorrhagic strokes
- Hypertensive heart disease
- Atrial fibrillation and flutter
- Aortic aneurysm
- Peripheral vascular disease
- Other cardiovascular and circulatory diseases

Orthopaedic

- Hip fracture
- Non-hip fracture

Other

- Diabetes mellitus type 2
- Peptic ulcer disease
- Rheumatoid arthritis
- Fire injuries

Cancer

- Lip and oral cavity cancer
- Nasopharynx cancer
- Cancer of nasal cavity and accessory sinuses
- Oesophageal cancer
- Stomach cancer
- Colon and rectum cancer
- Pancreatic cancer
- Larynx cancer
- Kidney cancer
- Bladder cancer
- Acute myeloid leukaemia
- Cervical cancer
- Liver cancer

Orthoptic

- Cataracts
- Macular degeneration

Respiratory

- Interstitial lung disease and pulmonary sarcoidosis
- Other chronic respiratory diseases
- Influenza and pneumonia
- Asthma in adolescents (for adults, smoking is only causally linked to exacerbation of existing asthma resulting in increased risk of hospitalisation and death)
- Tuberculosis

Reproduction and birth

- Reduced fertility in women
- Erectile dysfunction
- Antepartum haemorrhage
- Premature rupture of membranes
- Ectopic pregnancy
- Stillbirth

✓ Achieve your purpose of prioritising those with, or at risk of, poorer health²

While 16.9% of Victorian adults continue to be current smokers,³ those with, or at risk of, poorer health have much higher rates of tobacco use:

- **21.2%** of Australians 14 years and over in the most disadvantaged SEIFA quintile were **current** smokers in 2019 (compared to **8.1%** in the least disadvantaged SEIFA quintile).⁴
- **32.5%** of Australians that were unable to work were **current** smokers in 2019.⁴
- **29.9%** of single Australians with dependent children were **current** smokers in 2019.⁴
- **37%** of Aboriginal and Torres Strait Islander Australians aged 15 and over smoked **daily** in 2018–19.⁵
- **24.2%** of Australian adults diagnosed or treated for a mental illness in the last year were current smokers in 2019 (compared to **12.9%** who were not diagnosed or treated).⁴
- **23%** of homosexual/bisexual Australians were **current** smokers in 2019.⁴
- **77%** of homeless people and **93%** of “street” homeless people (those usually dwelling on streets, in parks, in derelict buildings or other temporary shelters) have an **elevated prevalence** of smoking.⁶
- Some **culturally and linguistically diverse** communities⁷
 - Those residing in metropolitan Melbourne
 - Those that are younger
 - Arabic-speaking people from a Lebanese background
 - Mandarin-speaking males from a Chinese background
 - Males from a Vietnamese background
 - Hindi-speaking males from an Indian background
 - Smokers of shisha (particularly in Hindi and Arabic communities).

✓ Reduce the COVID-19 risks and consequences exacerbated by smoking and vaping

- Smoking has **substantial adverse effects** on the immune system.⁸
 - Smoking disrupts the normal functioning of the immune system that fights infection in the **respiratory tract**.⁸
 - Among younger people (less than 69 years) current smokers were **nearly twice as likely** as never smokers to become infected with the COVID-19 virus.⁸
 - Among older people (69+ years), smokers were **more likely** to die from COVID-19 than non-smokers.⁸
 - Smokers have an **increased risk of contracting** COVID-19 and an **increased risk of needing to be hospitalised** with COVID-19.⁸
 - Current smokers with COPD, cancer, diabetes and those needing urgent surgery for hip fractures are at **greater risk of complications and death** from COVID-19.⁸
 - Vaping has been linked to substantially **increased risk of COVID-19** in teens and young adults⁹ and to more severe COVID-19 symptoms.¹⁰
 - Preliminary evidence* hints at the possibility that smoking may **reduce the effectiveness** of COVID-19 vaccination.
- *An Italian study found a lower level of antibodies in smokers compared to non-smokers 1–4 weeks after the second dose of the Pfizer vaccine.¹¹

✓ Act early on concerns that the increasing use of e-cigarettes will lead to increases in smoking in young people

- Growing evidence suggests a possible relationship between e-cigarette use and the **uptake and escalation** of smoking in young people.¹²
- The use of e-cigarettes is growing in Australia with the **highest proportion** of ever users among current smokers and those aged 18–24 years.¹³
- For non-smokers aged 18–24 years, 4.9% used them in 2013, 13.6% in 2016 and **19.6%** in 2019.¹³
- In an Australian survey of school students, around **14%** of students aged **12–17 years** had used an e-cigarette (32% of these in the past month).¹⁴
- **Almost half (48%)** of school students who vaped had never smoked tobacco before trying an e-cigarette.¹⁴
- Around a quarter of these students who had used e-cigarettes before ever smoking, reported later **trying tobacco cigarettes**.¹⁴

✓ Improve mental health and wellbeing

- Smoking has been associated with the development of increasing **social isolation** and **loneliness** in older adults aged 50 years and over.¹⁵
- Quitting smoking for at least six weeks **improves mental health**, mood, and quality of life, and **reduces** depression, anxiety and stress, both among the general population and among people with a psychiatric disorder.¹⁵
- Quitting smoking for at least six weeks has the **same or larger effect** as that of antidepressant treatment for mood and anxiety disorders.¹⁶

✓ Tackle poverty in applying the social model of health²

- Due to increases in the tax on tobacco, the price of cigarettes has **increased substantially** with a ten-fold increase in cost between 1980 and 2020.¹⁷
- Multiple studies have found associations between **smoking, financial stress, and housing insecurity**.¹⁸
- Families where one or more parent uses tobacco not only suffer **more** immediate **financial stress**, but also less long-term financial security and a **greater** likelihood of **poverty**.¹⁸
- The lowest-income households' expenditure on tobacco products as a proportion of total household weekly expenditure has been shown to be **over double** that of the highest income households.¹⁹
- If smokers quit, their chances of experiencing financial stress **reduce substantially** when compared to those of continuing smokers.¹⁹

✓ Reduce exposure to the substantial adverse effects of second- and third-hand smoke on non-smokers and children

- **Adverse health outcomes** caused by secondhand smoke (cigarette smoke lingering in the air inhaled by those in the vicinity) include:
 - Low birthweight
 - Sudden Infant Death Syndrome (SIDS)
 - Asthma (children)
 - Lower respiratory illness (children)
 - Otitis media (children)
 - Oro-facial clefts
 - Lung cancer
 - Ischaemic heart disease
 - Cerebrovascular disease.¹
- There is also **emerging evidence** that secondhand smoke exposure:
 - increases the risk of developing type 2 diabetes mellitus²⁰
 - is associated with depressive symptoms, psychological distress and stress, including in the postpartum period²¹
 - has an impact on cognition and behaviour, including higher likelihood of childhood conduct problems and learning difficulties.²²
- Children from disadvantaged families are **far more likely** to be exposed to secondhand smoke at home.²³
- Emerging evidence indicates that thirdhand smoke (toxic residue from tobacco smoke that settles on surfaces that is inhaled, ingested or absorbed through skin) **increases the risk** of adverse health outcomes such as hyperactive behaviour in children.²⁴
- Young children are **more likely to be at risk** of thirdhand smoke exposure due to more time spent indoors, more interaction with contaminated surfaces and a tendency to put objects in their mouth.²⁴

✓ Facilitate and strengthen collaborative local and state-wide efforts to tackle tobacco-related harm

- In developing municipal public health and wellbeing plans (MPHWP), local councils are required by legislation to have regard for the Victorian Public Health and Wellbeing Plan.²⁵
- The Victorian Public Health and Wellbeing Plan 2019–23 identifies **reducing tobacco-related harm** as one of 10 health priorities and one of a subset of four priorities for **particular focus**.²⁵
- Local councils are encouraged to concentrate MPHWP on the four focus areas.²⁵
- Community Health – Health Promotion teams are required to collaborate with local partners and align prevention and health promotion efforts with MPHWP to **maximise impact**.²⁶

✓ Benefit from the support, guidance and opportunities offered by Quit Victoria to Community Health – Health Promotion teams

- In 2022, Quit is **expanding** its **support and guidance** of Community Health – Health Promotion teams to build the capacity of the sector to reduce tobacco-related harm.
- Quit will establish a Resource Hub on its website providing offerings **tailored** to the work practice and circumstances of Community Health - Health Promotion teams including:
 - practice guides and frameworks
 - social marketing toolkits
 - professional development
 - interactive peer support opportunities
 - case studies
 - tools and templates
 - resources.
- Community Health – Health Promotion teams will have the opportunity to **sign up** to a Community of Practice and to receive this enhanced support and guidance.
- Quit will **showcase and enable sharing** of the initiatives and work practice of Community Health – Health Promotion teams in acting to reduce tobacco-related harm.
- Quit will invite Community Health Service senior leaders to provide a **summary of why** they have chosen to **prioritise** a focus on **reducing tobacco-related harm** which will be published on Quit's website and shared with key stakeholders.

For further information contact quit@cancervic.org.au.

References

1. Whetton S, Tait RJ, Scollo M, Banks E, Chapman J, Dey T, et al. Identifying the social costs of tobacco use to Australia in 2015/16. Perth: The National Drug Research Institute at Curtin University; 2019. Available from: <https://ndri.curtin.edu.au/NDRI/media/documents/publications/T273.pdf>
2. Department of Health. Community Health Services [internet]. Melbourne: Department of Health; 2021 [updated 2021 Oct 6; cited 2021 Feb 7]. Available from: <https://www.health.vic.gov.au/community-health/community-health-services>
3. Victorian Agency for Health Information. Victorian Population Health Survey 2019: summary of results. Melbourne: State of Victoria; 2021. Available from: <https://www.bettersafecare.vic.gov.au/sites/default/files/2021-05/Victorian%20Population%20Health%20Survey%202019%20-%20final.pdf>
4. Australian Institute of Health and Welfare. National Drug Strategy Household Survey 2019. Drug Statistics series no. 32. Cat. no. PHE 270. Canberra: AIHW; 2020. Available from: <https://www.aihw.gov.au/getmedia/3564474e-f7ad-461c-b918-7f8de03d1294/aihw-phe-270-NDSHS-2019.pdf.aspx?inline=true>
5. Australian Bureau of Statistics. 4715.0 - National Aboriginal and Torres Strait Islander Health Survey, 2018-19. Canberra: ABS; 2019. Available from: [https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4715.0Main%20Features152018-19?opendocument&tabname=Summary&prodno=4715.0&issue=2018-19&num=&view="](https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4715.0Main%20Features152018-19?opendocument&tabname=Summary&prodno=4715.0&issue=2018-19&num=&view=)
6. Kermod M, Crofts N, Miller P, Speed B, Streeton J. Health indicators and risks among people experiencing homelessness in Melbourne, 1995-1996. Aust NZ J Public Health. 1998;22:464-70. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-842X.1998.tb01415.x?sid=nlm%3Apubmed>
7. Mitsopoulos E, Durkin S. Research insights to inform the design of Quit's program of work with CALD communities. Melbourne: Cancer Council Victoria, Centre for Behavioural Research in Cancer; 2021 (unpublished report).
8. Winnall WR, Bellow B, Greenhalgh EM, Winstanley MH. 3.9 Increased susceptibility to infection in smokers. In Greenhalgh EM, Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2021. Available from: <http://www.tobaccoaustralia.org.au/3-9-increased-susceptibility-to-infection-in-smoke>
9. Gaiha SM, Cheng J, Halpern-Felsher B. Association Between Youth Smoking, Electronic Cigarette Use, and COVID-19. J Adolesc Health. 2020;67(4):519-523. Available from: [https://www.jahonline.org/article/S1054-139X\(20\)30399-2/pdf](https://www.jahonline.org/article/S1054-139X(20)30399-2/pdf)
10. McFadden DD, Bornstein SL, Vassallo R, Salonen BR, Bhuiyan NM, Schroeder DR, et al. Symptoms COVID 19 Positive Vapers Compared to COVID 19 Positive Non-vapers. J Prim Care Community Health. 2022;13:1-10. Available from: <https://journals.sagepub.com/doi/full/10.1177/21501319211062672>
11. Watanabe M, Balena A, Tuccinardi D, Tozzi R, Risi R, Masi D, et al. Central obesity, smoking habit, and hypertension are associated with lower antibody titres in response to COVID-19 mRNA vaccine. Diabetes Metab Res Rev. 2022;38(1):e3465. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/dmrr.3465>
12. Greenhalgh EM, Scollo MM. 18B.6 Effects of e-cigarette use on smoking. In Greenhalgh EM, Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2021. Available from: <http://www.tobaccoaustralia.org.au/chapter-18-harm-reduction/indepth-18b-e-cigarettes>
13. Greenhalgh EM, Jenkins S, Scollo MM. 18B.3 Prevalence of e-cigarette use. In Greenhalgh EM, Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2021. Available from: <http://www.tobaccoaustralia.org.au/chapter-18-harm-reduction/indepth-18b-e-cigarettes>
14. Guerin N, White V. ASSAD 2017 Statistics & Trends: Australian Secondary Students' Use of Tobacco, Alcohol, Over-the-counter Drugs, and Illicit Substances. 2nd ed. Melbourne: Cancer Council Victoria; 2020. Available from: <https://www.health.gov.au/sites/default/files/documents/2020/07/secondary-school-students-use-of-tobacco-alcohol-and-other-drugs-in-2017.pdf>
15. Philip K, Bu F, Polkey MI, Brown J, Steptoe A, Hopkinson NS, et al. Relationship of smoking with current and future social isolation and loneliness: 12-year follow-up of older adults in England. Lancet. 2022;14:100302. Available from: [https://www.thelancet.com/journals/lanepi/article/PIIS2666-7762\(21\)00288-X/fulltext](https://www.thelancet.com/journals/lanepi/article/PIIS2666-7762(21)00288-X/fulltext)
16. Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P. Change in mental health after smoking cessation: systematic review and meta-analysis. Br Med J. 2014;348:g1151. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24524926>
17. Scollo M, Bayly M. 13.3 The price of tobacco products in Australia. In Greenhalgh EM, Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2021. Available from: <http://www.tobaccoaustralia.org.au/chapter-13-taxation/13-3-the-price-of-tobacco-products-in-australia>
18. Greenhalgh EM, Scollo MM, Pearce M. 9.5 Smoking and intergenerational poverty. In Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2016. Available from: <http://www.tobaccoaustralia.org.au/chapter-9-disadvantage/9-5-smoking-and-intergenerational-poverty>
19. Greenhalgh EM, Scollo MM, & Pearce M. 9.4 The relationship between tobacco smoking and financial stress. In Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2016. Available from: <http://www.tobaccoaustralia.org.au/chapter-9-disadvantage/9-4-the-relationship-between-tobacco-smoking-and-f>
20. Greenhalgh EM, Scollo MM, Pearce M. 9.1 Smoking prevalence and exposure to secondhand smoke among priority populations in Australia. In Greenhalgh EM, Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2021. Available from: <http://www.tobaccoaustralia.org.au/chapter-9-disadvantage/9-1-socioeconomic-position-and-disparities-in-toba>
21. Campbell MA, Ford C, Winstanley MH. 4.13 Secondhand smoke and type 2 diabetes mellitus. In Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2017. Available from <http://www.tobaccoaustralia.org.au/chapter-4-secondhand/4-13-secondhand-smoke-and-diabetes>
22. Campbell MA, Ford C, Winstanley MH. 4.14 Secondhand smoke and mental health. In Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2017. Available from <http://www.tobaccoaustralia.org.au/chapter-4-secondhand/4-14-secondhand-smoke-and-mental-health>
23. Campbell MA, Winnall WR, Ford C, Winstanley MH. 4.17 Health effects of secondhand smoke for infants and children. In Greenhalgh EM, Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2021. Available from <http://www.tobaccoaustralia.org.au/chapter-4-secondhand/4-17-health-effects-of-secondhand-smoke-for-infants>
24. Campbell MA, Ford C, Winstanley MH. 4.3 Thirdhand smoke. In Scollo MM and Winstanley MH [editors]. Tobacco in Australia: Facts and issues. Melbourne: Cancer Council Victoria; 2017. Available from <http://www.tobaccoaustralia.org.au/chapter-4-secondhand/4-3-thirdhand-smoke>
25. Department of Health and Human Services. Municipal public health and wellbeing planning 2021-2025: Advice Note 1. Melbourne: Department of Health and Human Services; 2020. Available from: https://www.mav.asn.au/__data/assets/pdf_file/0019/26254/MPHWP-2021-2025-Advice-Note-1-August2020.pdf
26. Department of Health. Community Health – Health Promotion draft program guidelines 2021-2025. Melbourne: Department of Health; 2021.